medRxiv preprint doi: https://doi.org/10.1101/2020.12.02.20240648; this version posted December 4, 2020. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.
It is made available under a CC-BY 4.0 International license .

The effectiveness of population-wide, rapid antigen test based screening in reducing SARS-CoV-2 infection prevalence in Slovakia

Authors

Martin Pavelka [1,2,3], Kevin Van-Zandvoort [4,5], Sam Abbott [4,5], Katharine Sherratt [4,5], Marek Majdan [7], CMMID COVID-19 working group [5], Inštitút Zdravotných Analýz [6], Pavol Jarčuška [7], Marek Krajčí [1+], Stefan Flasche [4,5*], Sebastian Funk [4,5*]

[1] Slovak Ministry of Health, Bratislava, Slovakia
[2] Inštitút Zdravotných Analýz, Bratislava, Slovakia
[3] Department of Public Health and Policy, London School of Hygiene & Tropical Medicine, London, UK
[4] Department for Infectious Disease Epidemiology, London School of Hygiene & Tropical Medicine, London, UK
[5] Centre for Mathematical Modelling of Infectious Diseases, London School of Hygiene & Tropical Medicine, London, UK
[6] Inštitút Zdravotných Analýz (Institute of Health Analyses), Bratislava, Slovakia
[7] Faculty of Medicine, Pavol Jozef Šafárik University, Košice, Slovakia
[7] Institute for Global Health and Epidemiology, Faculty of Health Sciences and Social Work, Trnava University, Trnava, Slovakia

[+] on behalf of all Slovak medical staff who have helped in the mass testing campaigns
* authors contributed equally, order decided by coin toss
Address correspondence to: Martin Pavelka <martin.pavelka@health.gov.sk>

Abstract

Non-pharmaceutical interventions have been extensively used worldwide to limit the transmission of SARS-CoV-2, but they also place an enormous social and economic burden on populations . We report the results of recent mass testing for SARS-CoV-2 in Slovakia where rapid antigen tests were used to screen the whole population and to isolate infectious cases together with their household members. Prevalence of detected infections decreased by 58% (95% CI: 57-58%) within one week in the 45 counties that were subject to two rounds of mass testing. Adjusting for geographical clustering and differences in attendance rates and the epidemiological situation at the time of the first round, this changed to 61% (95% CI: 50-70%). Adjusting for an estimated growth rate in infections of 4.4% (1.1-6.9%) per day in the week preceding the mass testing campaign and the corresponding expected growth in infection prevalence, the estimated decrease in prevalence compared to a scenario of unmitigated growth was 70% (67-73%). Using a microsimulation model we find that this decrease can not be explained solely by infection control measures that were introduced in the weeks preceding the intervention, but requires the additional impact of isolation as well as quarantine of household members of those testing positive during the mass testing campaign.

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

medRxiv preprint doi: https://doi.org/10.1101/2020.12.02.20240648; this version posted December 4, 2020. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.
It is made available under a CC-BY 4.0 International license .

Introduction

Non-pharmaceutical interventions have been extensively used worldwide to limit the transmission of SARS-CoV-2 (*1–4*). These have included travel restrictions, mandating face masks, closure of schools and non-essential businesses, and nationwide stay-at-home orders. While all the measures were aimed at mitigating ill-health due to COVID-19 (*3*, *5*) they also place an unprecedented economic and social burden on people (*6–9*), the majority uninfected. Testing of reported symptomatic cases and tracing their contacts aims to provide a more targeted measure but in many settings has proven insufficient for containing transmission (*10*, *11*).

Mass testing campaigns are an alternative way to identify infectious individuals and allow targeting of interventions without much added burden to those uninfectious (*12*). However, they have been limited until recently by the dependence on Polymerase Chain Reaction (PCR) for the diagnosis of a SARS-CoV-2 infection. While laboratory capacities have been upscaled in record time, PCR testing remains expensive and can seldom achieve a turnaround time of less than one day (*13*, *14*). In comparison, recently developed rapid antigen tests are cheap and can be quickly produced in large quantities offering results on site in 15-30 mins without the need for a laboratory. They are less sensitive in detecting infections with low viral load but have been found to detect the vast majority of infectious infections, and hence may make mass testing a viable part of the portfolio of non-pharmaceutical interventions (*15–17*).

In October 2020, Slovakia became the first country in the world to use rapid antigen tests in a campaign targeting the whole population in order to identify infections at scale, rapidly reduce transmission and allow quicker easing of lockdown measures(*18*). A pilot took place between 23 and 25 October in the four most affected counties, followed by a round of national mass testing on 31 October and 1 November (henceforth: round 1). High prevalence counties were again targeted with a subsequent round on 7 and 8 November (round 2).

We evaluated the impact of mass testing in Slovakia, in combination with other measures put in place around the time, by comparing infection prevalence in each round of testing.

Results

In total, 5,276,832 rapid antigen tests were used in the mass testing campaigns, with 65% of the respective populations tested in the pilot, 66% in mass testing round 1 and 62% round 2. This corresponded to 87%, 83% and 84% of the age-eligible population in each round, respectively, and does not include another 534,300 tests that were conducted through additional testing sites for medical, military and governmental personnel and not included in geographical county data.

A total of 50,466 tests indicated the presence of a currently infectious SARS-CoV-2 infection. The proportion of positive tests was 3.91% (range across counties: 3.12 to 4.84%) in the pilot, 1.01% (range: 0.13-3.22%) in round 1 and 0.62% (range: 0.28-1.65%) in

medRxiv preprint doi: https://doi.org/10.1101/2020.12.02.20240648; this version posted December 4, 2020. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.
It is made available under a CC-BY 4.0 International license .

round 2 (Figure 2C and D). We estimate that with 95% certainty the specificity of the SD Biosensor Standard Q antigen test was exceeding 99.85%.

In the four counties where the pilot was conducted, prevalence decreased by 56% (95% Confidence Interval, CI: 54-58%) between the pilot and round 1 of the mass testing campaign and a further 60% (95% CI: 56-63%) between rounds 1 and 2, totalling a decrease of 82% (95% CI: 81-83%) over two weeks. There was little heterogeneity between counties (Figure 2B).

Among the 45 counties that were included in round 2 of the mass testing campaign, infection prevalence decreased by 58% (95% CI: 57-58%) in the crude analysis and by 61% (95% CI: 50-70%) if adjusted for differences in region, attendance rates, reproduction number and round 1 prevalence. The estimated reduction varied by county from 29% in county Považská Bystrica to 79% in county Medzilaborce but with little regional differences (Figure 2A). Neither region, attendance rates, prevalence in round 1 or the estimated growth rate prior to mass testing were found to be significantly associated with county specific reductions.

At the time of round 1 of the mass testing campaign incidence of confirmed cases was rising in non-pilot counties with an estimated infection growth rate of 4.4% (1.1%-6.9%) per day. When adjusting for this growth trend, we estimated a self-adjusted prevalence ratio (saPR) of 0.30 (0.27-0.33) . In the pilot counties, reported infection incidence showed signs of levelling in the week before the mass testing campaign with an estimated infection growth rate of 1.3% (-7.4-7.8%), yielding a respective saPR of 0.31 (0.26-0.33).

In a microsimulation model, only the scenario that assumed a substantial impact of both the lockdown and the mass testing was able to generate reductions in test positivity rates between testing rounds that were similar to those observed (Figure 3). The requirement for quarantine for the whole household following a positive test was essential for the effect of mass testing; predicted prevalence ratio between the first two testing rounds of 0.41 (0.38-0.45) with and 0.90 (0.84-0.96) without household quarantine.

Discussion

The reduction in prevalence achieved in Slovakia through a combination of restrictions on movement and the first ever large scale rapid antigen mass testing is striking, with reductions of over 50% achieved within a week between two rounds of testing. While we could not with certainty disentangle the effects, simulations from a mathematical model suggested that both the restrictions and mass testing likely contributed substantially to the observed impact and that quarantining of household contacts was a crucial contribution to the effectiveness of mass testing.

Potentially large numbers of false positive tests have been a point of criticism for mass testing campaigns. While multiple studies have found high specificity of the Biosensor test kit they are not powered to exclude specificity levels that on population level would yield an overwhelming amount of false positives (*15*, *22*). We show that indeed specificity is very likely exceeding 99.85% and therefore not of major concern in this study.

While we observed a dramatic reduction in test positivity between mass testing campaigns, the observed change in daily case incidence reported through standard

medRxiv preprint doi: https://doi.org/10.1101/2020.12.02.20240648; this version posted December 4, 2020. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.
It is made available under a CC-BY 4.0 International license .

surveillance was not the rapid collapse in test-positive cases that would correspond to the drastic reductions in prevalence. This may be due to a variety of reasons. Foremost, national mass testing campaigns are likely to have a major disruptive effect on passive syndromic surveillance. In addition, starting mid-September the incidence surveillance has been operating at capacity with long waiting lists for testing and stricter eligibility criteria, which in the post mass testing period reduced substantially, and hence may have artificially reduced the observable change in such data. In contrast, data on hospital bed occupancy shows sudden flattening from mid-November suggesting a sharp decrease in new admissions consistent with a sizable reduction in new infections at the time of the mass testing campaigns (Figure S6).

The most important limitation of this observational study is that we were unable to clearly distinguish the effect of the mass testing campaigns from that of the other non-pharmaceutical interventions introduced at a similar time, that have led to a reduction in contacts and mobility, albeit much less than during the Spring lockdown (Figure S4). We are unaware of any other context in which a COVID-19 intervention has resulted in a 60% decline in infection prevalence within one week (or 80% in two weeks), particularly while primary schools and workplaces were mostly open. This would suggest that indeed a large share of the impact can be attributed to the mass testing campaigns. Similarly, our analysis using mathematical modelling suggests that even with what would be considered as one of the most impactful lockdowns observed so far, it would be impossible to replicate such rapid drop in test positivity without a substantial contribution from the mass testing campaign.

The need to mobilise sufficient medical personnel to conduct the nasopharyngeal swabs could be a major obstacle to countries. Other rapid antigen tests kits are available that have achieved similarly high sensitivity in detecting likely infectious infections in lab conditions but are also licensed for use with nasal swabs (*32*, *33*). Nasal swabs can be self-administered and therefore reduce demand on trained personnel and transmission risk in the process of sample collection or even may enable testing at home. However, these benefits have to be carefully weighed against the potential loss of sensitivity if self administered (*34*).

In conclusion, the combination of nationwide restrictions and mass testing with quarantining of household contacts of test positives rapidly reduced the prevalence of infectious residents in Slovakia. While impossible to disentangle the precise contribution of control measures and mass testing, the latter is likely to have had a substantial effect in curbing the pandemic in Slovakia and may provide a key tool in the containment of SARS-CoV-2.

Material and Methods

Study population

Slovakia is a country with a population of 5.5 million, consisting of 79 counties grouped into 8 administrative regions. Slovak residents aged between 10 and 65 years and older adults in employment were eligible for mass testing (about 4 million people). Those

medRxiv preprint doi: https://doi.org/10.1101/2020.12.02.20240648; this version posted December 4, 2020. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.
It is made available under a CC-BY 4.0 International license .

quarantining at the time or who had recovered from COVID-19 in the past three months were excluded.

The pilot was conducted in three counties in the Orava subregion (Námestovo, Tvrdošín, Dolný Kubín) and Bardejov county, which had the highest infection incidence at the time. The first round of mass testing was conducted nation-wide and the second round of mass testing was restricted to 45 counties, mostly in the northern part of Slovakia, with infection prevalence in the first mass testing round exceeding 7 per 1,000 tests.

Interventions

Slovakia implemented a series of infection control measures throughout October, which included closing schools for pupils aged 14 or above on 15 Oct and for pupils aged 10 and above on 26 October. They remained closed throughout the period of the mass testing campaigns and thereafter. Indoor gastronomy and indoor leisure activities were also restricted. Residents were further asked to limit their movement for one week between 24 October and 1 November only to: going to work, taking children to school, shopping for essential items and going for recreational walks (Figures 1 and S4). Although these rules were legally enforceable, Slovakia relied mostly on people's civil responsibility to adhere to restrictions.

On the days of mass testing, participants attended testing centres run by healthcare professionals, armed forces and volunteers. Overall, Slovakia deployed around twenty thousand medical staff and forty thousand non-medical personnel. Testing procedures followed as recommended by the manufacturer, with nasopharyngeal samples obtained by trained medical personnel using flexible, aluminum-shaft, calcium alginate swabs (*19*).

Testing was not obligatory, but residents who did not attend the mass testing were instructed to stay home for ten days or until the next round of mass testing. A medical certificate was issued to every participant confirming their infection status. A test-negative certificate was required by employers to enter workplaces. Various venues and public institutions inspected peoples' certificates at random. Private PCR tests were also accepted if no older than the most recent mass testing campaign. Citizens whose test results were positive were asked to enter a 10-day long quarantine together with all members of the same household and their self-traced contacts in the preceding two days in an attempt to reduce secondary transmission.

Data

No participant information was collected during either of the mass-testing campaigns. However, information on the number of tests used as well as the number of positive tests has been tracked and made openly available by the Slovak Government (*18*). The SD Biosensor Standard Q antigen test that was used exclusively has high specificity, with point estimates typically in excess of 99.5%. Sensitivity exceeded 70% in most validation studies, and exceeded 90% among samples with a cycle threshold below 25, a threshold commonly associated with effective transmission (*15*, *20*–*22*).

medRxiv preprint doi: https://doi.org/10.1101/2020.12.02.20240648; this version posted December 4, 2020. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.
It is made available under a CC-BY 4.0 International license .

To assess trends in the local epidemiology of SARS-CoV we used routine syndromic and PCR confirmed surveillance for the daily incidence of infections as reported by the Slovak Ministry of Health (*23*).

Analyses

We calculated crude prevalence ratios (cPR) to estimate the change in test positivity between mass testing campaigns, including Wald-Normal confidence intervals. Binomial confidence intervals were calculated for prevalence estimates. Test positive rates provide a natural upper bound for false positive rates of a test. We thus estimated the minimum test specificity *ms* as the probability of observing a test positivity of at least 1-*ms* in at least one county, assuming the test positivity to be binomially distributed.

To explore heterogeneity between counties in the estimated reduction in test positivity in subsequent rounds of mass testing, we used a quasi-Poisson regression model. The number of positive tests in each county was modelled with a county specific intercept, an indicator variable for the round 2 of mass testing, and interactions of the latter with attendance rates in round 1, round 1 test positivity, the reproduction number leading up to round 1 and region as covariates as well as the log number of tests as an offset variable. The three continuous variable interaction terms were centered and standardised (see supplement).

We used the EpiNow2 model (*24*, *25*) for the calculation of trends in local epidemiology prior to mass testing based on routinely reported infection incidence. EpiNow2 uses observed delay distributions in combination with a renewal equation model to probabilistically infer the infection date for each reported case as well as the population-wide time varying reproduction number (*26–28*) , allowing a smoothed extrapolation of infection incidence and prevalence and extrapolation beyond the observed study period under an assumption of no change. We define the self-adjusted prevalence ratio (saPR) as the cPR divided by the prevalence ratio at the times of round 2 vs round 1 as estimated through EpiNow2. The saPR is an estimate for the effect of the intervention that takes into account that infection prevalence would have changed in the time between observations (see supplement).

To explore scenarios for the relative effect of mass testing and lockdown we used a microsimulation model. We focused on three scenarios in which mass testing takes place, i) an epidemic growth rate of R_e=1.4 (as in early October) that is unchanged by lockdown measures, ii) a reduced growth rate of R_e=1 from 15th October (similar to many parts of Europe in the weeks following autumn lockdowns) and iii) the growth rate reduced to R_e=0.6 from 15th October (the smallest observed reproduction number nationally during the COVID-19 pandemic) but no effect of mass testing. A detailed model description is provided in the supplementary material, but in brief: Individuals are grouped in households according to Slovak census data (*29*), and make contact with individuals outside their household at age-specific rates(*30*). To account for social distance measures, we assumed absence of at-school contacts for children 10 years and over, and that contacts at work and contacts not at the home, school, or workplace, were reduced by 25% and 75% from pre-epidemic levels, respectively. We simulated infections among 78,000 susceptible individuals, representative of the population size of a typical pilot county. When infection prevalence reached 3.2% (approximating a typical observed

medRxiv preprint doi: https://doi.org/10.1101/2020.12.02.20240648; this version posted December 4, 2020. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.
It is made available under a CC-BY 4.0 International license .

prevalence during the testing pilot), up to 3 rounds of weekly mass testing were initiated and the week before that restrictions equivalent to those enacted in Slovakia were implemented. In the model, we assumed perfect test sensitivity for detection of currently infectious infections, specificity, and compliance with quarantine. Observed test attendance rates were used assuming that individuals in quarantine did not attend mass-testing.

Open Access

Daily incidence of positive COVID-19 test reports and the results of the mass testing are available through governmental websites (*18*, *23*). All analyses were conducted in R (*31*) and can be found at www.github.com/sbfnk/covid19.slovakia.mass.testing (data analyses) and https://github.com/kevinvzandvoort/covid_svk (simulation model).

Funding

Martin Pavelka is employed by the Slovak Ministry of Health. Marek Krajčí is a medical doctor, member of the Slovak government and Slovak Minister of Health. Stefan Flasche is supported by a Sir Henry Dale Fellowship jointly funded by the Wellcome Trust and the Royal Society (Grant number 208812/Z/17/Z). Sebastian Funk, Sam Abbott and Katharine Sherratt are supported by the Wellcome Trust (210758/Z/18/Z). Kevin van Zandvoort is supported by Elrha's Research for Health in Humanitarian Crises (R2HC) Programme, which aims to improve health outcomes by strengthening the evidence base for public health interventions in humanitarian crises. The R2HC programme is funded by the UK Government (DFID), the Wellcome Trust, and the UK National Institute for Health Research (NIHR).

The following funding sources are acknowledged as providing funding for the working group authors. BBSRC LIDP (BB/M009513/1: DS). This research was partly funded by the Bill & Melinda Gates Foundation (INV-001754: MQ; INV-003174: KP, MJ, YL; NTD Modelling Consortium OPP1184344: CABP, GFM; OPP1180644: SRP; OPP1183986: ESN). BMGF (OPP1157270: KA). DFID/Wellcome Trust (Epidemic Preparedness Coronavirus research programme 221303/Z/20/Z: CABP). EDCTP2 (RIA2020EF-2983-CSIGN: HPG). ERC Starting Grant (#757699: MQ). This project has received funding from the European Union's Horizon 2020 research and innovation programme - project EpiPose (101003688: KP, MJ, PK, RCB, WJE, YL). This research was partly funded by the Global Challenges Research Fund (GCRF) project 'RECAP' managed through RCUK and ESRC (ES/P010873/1: AG, CIJ, TJ). HDR UK (MR/S003975/1: RME). MRC (MR/N013638/1: NRW). Nakajima Foundation (AE). This research was partly funded by the National Institute for Health Research (NIHR) using UK aid from the UK Government to support global health research. The views expressed in this publication are those of the author(s) and not necessarily those of the NIHR or the UK Department of Health and Social Care (16/136/46: BJQ; 16/137/109: BJQ, FYS, MJ, YL; Health Protection Research Unit for Immunisation NIHR200929: NGD; Health Protection Research Unit for Modelling Methodology HPRU-2012-10096: TJ; NIHR200908: RME; NIHR200929: FGS, MJ; PR-OD-1017-20002: AR, WJE). Royal Society (Dorothy Hodgkin Fellowship: RL; RP\EA\180004: PK). UK DHSC/UK Aid/NIHR (PR-OD-1017-20001: HPG). UK

medRxiv preprint doi: https://doi.org/10.1101/2020.12.02.20240648; this version posted December 4, 2020. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.
It is made available under a CC-BY 4.0 International license .

MRC (MC_PC_19065 - Covid 19: Understanding the dynamics and drivers of the COVID-19 epidemic using real-time outbreak analytics: AG, NGD, RME, SC, TJ, WJE, YL; MR/P014658/1: GMK). Authors of this research receive funding from UK Public Health Rapid Support Team funded by the United Kingdom Department of Health and Social Care (TJ). Wellcome Trust (206250/Z/17/Z: AJK, TWR; 206471/Z/17/Z: OJB; 208812/Z/17/Z: SC; 210758/Z/18/Z: JDM, JH, NIB, SA, SRM). No funding (AMF, AS, CJVA, DCT, JW, KEA, YWDC).

Conflicts of interest

All authors declare that they have no conflicts of interest

Acknowledgements

We would like to thank all healthcare workers, Slovak armed forces and countless volunteers who helped with the execution of the mass testing campaign. Lastly we would like to thank all participants who contributed their time to help curb the Pandemic and particularly those who had to quarantine as a result of their or their household's or contact's test result.

The following authors were part of the Centre for Mathematical Modelling of Infectious Disease COVID-19 Working Group. Each contributed in processing, cleaning and interpretation of data, interpreted findings, contributed to the manuscript, and approved the work for publication: Gwenan M Knight, Naomi R Waterlow, Carl A B Pearson, Fiona Yueqian Sun, Simon R Procter, Alicia Showering, Rosalind M Eggo, Yung-Wai Desmond Chan, Emily S Nightingale, David Simons, Oliver Brady, Billy J Quilty, Petra Klepac, Amy Gimma, Hamish P Gibbs, W John Edmunds, Adam J Kucharski, Sam Abbott, Jack Williams, Kiesha Prem, Rosanna C Barnard, Thibaut Jombart, Graham Medley, Katherine E. Atkins, Samuel Clifford, Nicholas G. Davies, Kaja Abbas, Mark Jit, Timothy W Russell, Frank G Sandmann, Damien C Tully, James D Munday, Anna M Foss, Alicia Rosello, Sophie R Meakin, Joel Hellewell, C Julian Villabona-Arenas, Christopher I Jarvis, Rachel Lowe, Akira Endo, Matthew Quaife, Nikos I Bosse, Yang Liu.

Literature

1. Guidelines for the implementation of non-pharmaceutical interventions against COVID-19. *Eur. Cent. Dis. Prev. Control* (2020), (available at https://www.ecdc.europa.eu/en/publications-data/covid-19-guidelines-non-pharmaceutical-interventions).
2. B. J. Cowling, S. T. Ali, T. W. Y. Ng, T. K. Tsang, J. C. M. Li, M. W. Fong, Q. Liao, M. Y. Kwan, S. L. Lee, S. S. Chiu, J. T. Wu, P. Wu, G. M. Leung, Impact assessment of non-pharmaceutical interventions against coronavirus disease 2019 and influenza in Hong Kong: an observational study. *Lancet Public Health*. **5**, e279–e288 (2020).
3. N. G. Davies, A. J. Kucharski, R. M. Eggo, A. Gimma, W. J. Edmunds, T. Jombart, K. O'Reilly, A. Endo, J. Hellewell, E. S. Nightingale, B. J. Quilty, C. I. Jarvis, T. W. Russell, P. Klepac, N. I. Bosse, S. Funk, S. Abbott, G. F. Medley, H. Gibbs, C. A. B. Pearson, S. Flasche, M. Jit, S. Clifford, K. Prem, C. Diamond, J. Emery, A. K. Deol, S. R. Procter, K. van Zandvoort, Y. F. Sun, J. D. Munday, A. Rosello, M. Auzenbergs, G. Knight, R. M. G. J. Houben, Y. Liu, Effects of non-pharmaceutical interventions on COVID-19 cases, deaths, and demand for hospital services in the UK: a modelling study. *Lancet Public Health*. **0** (2020), doi:10.1016/S2468-2667(20)30133-X.

medRxiv preprint doi: https://doi.org/10.1101/2020.12.02.20240648; this version posted December 4, 2020. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.
It is made available under a CC-BY 4.0 International license .

4. J. M. Brauner, S. Mindermann, M. Sharma, D. Johnston, J. Salvatier, T. Gavenčiak, A. B. Stephenson, G. Leech, G. Altman, V. Mikulik, A. J. Norman, J. T. Monrad, T. Besiroglu, H. Ge, M. A. Hartwick, Y. W. Teh, L. Chindelevitch, Y. Gal, J. Kulveit, *medRxiv*, in press, doi:10.1101/2020.05.28.20116129.
5. N. Ferguson, D. Laydon, G. Nedjati Gilani, N. Imai, K. Ainslie, M. Baguelin, S. Bhatia, A. Boonyasiri, Z. Cucunuba Perez, G. Cuomo-Dannenburg, A. Dighe, I. Dorigatti, H. Fu, K. Gaythorpe, W. Green, A. Hamlet, W. Hinsley, L. Okell, S. Van Elsland, H. Thompson, R. Verity, E. Volz, H. Wang, Y. Wang, P. Walker, P. Winskill, C. Whittaker, C. Donnelly, S. Riley, A. Ghani, "Report 9: Impact of non-pharmaceutical interventions (NPIs) to reduce COVID19 mortality and healthcare demand" (Imperial College London, 2020), , doi:10.25561/77482.
6. Spotlight on the SDGs | Digital library: Publications. *UN Women*, (available at https://www.unwomen.org/en/digital-library/publications/2018/4/spotlight-on-the-sdgs).
7. Everyone Included: Social Impact of COVID-19 | DISD, (available at https://www.un.org/development/desa/dspd/everyone-included-covid-19.html).
8. Socio-economic impact of COVID-19. *UNDP*, (available at https://www.undp.org/content/undp/en/home/coronavirus/socio-economic-impact-of-covid-19.html).
9. M. Polyakova, G. Kocks, V. Udalova, A. Finkelstein, Initial economic damage from the COVID-19 pandemic in the United States is more widespread across ages and geographies than initial mortality impacts. *Proc. Natl. Acad. Sci.* **117**, 27934–27939 (2020).
10. A. J. Kucharski, P. Klepac, A. J. K. Conlan, S. M. Kissler, M. L. Tang, H. Fry, J. R. Gog, W. J. Edmunds, CMMID COVID-19 working group, Effectiveness of isolation, testing, contact tracing, and physical distancing on reducing transmission of SARS-CoV-2 in different settings: a mathematical modelling study. *Lancet Infect. Dis.* **20**, 1151–1160 (2020).
11. S. Contreras, J. Dehning, M. Loidolt, F. P. Spitzner, J. H. Urrea-Quintero, S. B. Mohr, M. Wilczek, J. Zierenberg, M. Wibral, V. Priesemann, The challenges of containing SARS-CoV-2 via test-trace-and-isolate. *ArXiv200905732 Q-Bio* (2020) (available at http://arxiv.org/abs/2009.05732).
12. J. Peto, Covid-19 mass testing facilities could end the epidemic rapidly. *BMJ*. **368** (2020), doi:10.1136/bmj.m1163.
13. Weekly statistics for NHS Test and Trace (England) and coronavirus testing (UK): 5 November to 11 November. *GOV.UK*, (available at https://www.gov.uk/government/publications/nhs-test-and-trace-england-and-coronavirus-testing-uk-statistics-5-november-to-11-november/weekly-statistics-for-nhs-test-and-trace-england-and-coronavirus-testing-uk-5-november-to-11-november).
14. M. E. Kretzschmar, G. Rozhnova, M. C. J. Bootsma, M. van Boven, J. H. H. M. van de Wijgert, M. J. M. Bonten, Impact of delays on effectiveness of contact tracing strategies for COVID-19: a modelling study. *Lancet Public Health*. **5**, e452–e459 (2020).
15. Z. Iglói, J. Velzing, J. van Beek, D. van de Vijver, G. Aron, R. Ensing, K. Benschop, W. Han, T. Boelsums, M. Koopmans, C. Geurtsvankessel, R. Molenkamp, *medRxiv*, in press, doi:10.1101/2020.11.18.20234104.
16. Options for the use of rapid antigen tests for COVID-19 in the EU/EEA and the UK. *Eur. Cent. Dis. Prev. Control* (2020), (available at

medRxiv preprint doi: https://doi.org/10.1101/2020.12.02.20240648; this version posted December 4, 2020. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.
It is made available under a CC-BY 4.0 International license .

https://www.ecdc.europa.eu/en/publications-data/options-use-rapid-antigen-tests-covid-19-eueea-and-uk).

17. M. J. Mina, R. Parker, D. B. Larremore, *N. Engl. J. Med.*, in press, doi:10.1056/NEJMp2025631.
18. Som zodpovedny. *Som Zodp.*, (available at https://www.somzodpovedny.sk/).
19. Products - STANDARD Q COVID-19 Ag, (available at http://sdbiosensor.com/xe/product/7672).
20. L. J. Krüger, M. Gaeddert, L. Köppel, L. E. Brümmer, C. Gottschalk, I. B. Miranda, P. Schnitzler, H. G. Kräusslich, A. K. Lindner, O. Nikolai, F. P. Mockenhaupt, J. Seybold, V. M. Corman, C. Drosten, N. R. Pollock, A. I. Cubas-Atienzar, K. Kontogianni, A. Collins, A. H. Wright, B. Knorr, A. Welker, M. de Vos, J. A. Sacks, E. R. Adams, C. M. Denkinger, F. the S. Team, *medRxiv*, in press, doi:10.1101/2020.10.01.20203836.
21. F. Cerutti, E. Burdino, M. G. Milia, T. Allice, G. Gregori, B. Bruzzone, V. Ghisetti, Urgent need of rapid tests for SARS CoV-2 antigen detection: Evaluation of the SD-Biosensor antigen test for SARS-CoV-2. *J. Clin. Virol.* **132**, 104654 (2020).
22. L. Kaiser, I. Eckerle, M. Schibler, A. Berger, "Validation Report: SARS-CoV-2 Antigen Rapid Diagnostic Test," (available at https://www.hug.ch/sites/interhug/files/structures/laboratoire_de_virologie/documents/Centre_maladies_virales_infectieuses/ofsp_rdt_report_gcevd_27.10.2020.pdf).
23. Koronavírus a Slovensko. *Koronavírus Slov.*, (available at https://korona.gov.sk/).
24. S. Abbott, J. Hellewell, R. N. Thompson, K. Sherratt, H. P. Gibbs, N. I. Bosse, J. D. Munday, S. Meakin, E. L. Doughty, J. Y. Chun, Y.-W. D. Chan, F. Finger, P. Campbell, A. Endo, C. A. B. Pearson, A. Gimma, T. Russell, CMMID COVID modelling group, S. Flasche, A. J. Kucharski, R. M. Eggo, S. Funk, Estimating the time-varying reproduction number of SARS-CoV-2 using national and subnational case counts. *Wellcome Open Res.* **5**, 112 (2020).
25. *epiforecasts/EpiNow2* (epiforecasts, 2020; https://github.com/epiforecasts/EpiNow2).
26. J. Wallinga, P. Teunis, Different Epidemic Curves for Severe Acute Respiratory Syndrome Reveal Similar Impacts of Control Measures. *Am. J. Epidemiol.* **160**, 509–516 (2004).
27. R. N. Thompson, J. E. Stockwin, R. D. van Gaalen, J. A. Polonsky, Z. N. Kamvar, P. A. Demarsh, E. Dahlqwist, S. Li, E. Miguel, T. Jombart, J. Lessler, S. Cauchemez, A. Cori, Improved inference of time-varying reproduction numbers during infectious disease outbreaks. *Epidemics*. **29**, 100356 (2019).
28. S. Cauchemez, P.-Y. Boëlle, G. Thomas, A.-J. Valleron, Estimating in Real Time the Efficacy of Measures to Control Emerging Communicable Diseases. *Am. J. Epidemiol.* **164**, 591–597 (2006).
29. Statistics | Eurostat, (available at https://ec.europa.eu/eurostat/databrowser/view/cens_01rhsize/default/table?lang=en).
30. K. Prem, A. R. Cook, M. Jit, Projecting social contact matrices in 152 countries using contact surveys and demographic data. *PLOS Comput. Biol.* **13**, e1005697 (2017).
31. R Core Team, *R: A Language and Environment for Statistical Computing* (R Foundation for Statistical Computing, Vienna, Austria, 2019; https://www.R-project.org/).
32. Panbio COVID-19 Ag Rapid Test Device, (available at https://www.globalpointofcare.abbott/en/product-details/panbio-covid-19-ag-antigen-test.html).

medRxiv preprint doi: https://doi.org/10.1101/2020.12.02.20240648; this version posted December 4, 2020. The copyright holder for this preprint **(which was not certified by peer review)** is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.
It is made available under a CC-BY 4.0 International license .

33. Sofia SARS Antigen FIA | Quidel, (available at https://www.quidel.com/immunoassays/rapid-sars-tests/sofia-sars-antigen-fia).
34. E. Mahase, Covid-19: Innova lateral flow test is not fit for “test and release” strategy, say experts. *BMJ*. **371** (2020), doi:10.1136/bmj.m4469.

medRxiv preprint doi: https://doi.org/10.1101/2020.12.02.20240648; this version posted December 4, 2020. The copyright holder for this preprint **(which was not certified by peer review)** is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.
It is made available under a CC-BY 4.0 International license .

Tables and Figures

Table 1: Overview of county specific test numbers and reductions for the 79 counties in Slovakia. R: median estimate of the reproduction number on 22 October. %: proportion positive out of those attending mass testing.

				Round 1			Round 2		
County	Region	Population	R	Positive	Attendance	%	Positive	Attendance	%
Bánovce nad Bebravou	Trenčiansky	36282	1.4	457	23264	1.96	192	22248	0.86
Banská Bystrica	Banskobystrický	110828	1.2	687	64127	1.07	231	66544	0.35
Banská Štiavnica	Banskobystrický	16086	0.7	33	11725	0.28			
Bardejov	Prešovský	77771	0.7	740	44197	1.67	366	43983	0.83
Bratislava I	Bratislavský	44798	1.2	108	29047	0.37			
Bratislava II	Bratislavský	108139	1.2	345	80958	0.43			
Bratislava III	Bratislavský	61418	1.2	175	49788	0.35			
Bratislava IV	Bratislavský	93058	1.2	81	63857	0.13			
Bratislava V	Bratislavský	141259	1.2	268	68139	0.39			
Brezno	Banskobystrický	61450	1.4	450	37339	1.21	242	38515	0.63
Bytča	Žilinský	30917	1.6	328	21419	1.53	164	20931	0.78
Čadca	Žilinský	90080	1.0	1736	53907	3.22	506	52304	0.97
Detva	Banskobystrický	32051	1.3	211	19704	1.07	79	23255	0.34
Dolný Kubín	Žilinský	39456	1.0	345	24251	1.42	138	24170	0.57
Dunajská Streda	Trnavský	122358	1.3	840	87329	0.96	577	110083	0.52
Galanta	Trnavský	94076	1.3	349	71243	0.49			
Gelnica	Košický	31868	1.3	131	18331	0.71	72	19087	0.38
Hlohovec	Trnavský	45012	1.4	171	28892	0.59			
Humenné	Prešovský	61986	1.1	598	32962	1.81	197	32750	0.60
Ilava	Trenčiansky	59188	1.4	442	37604	1.18	291	35931	0.81
Kežmarok	Prešovský	75235	1.4	845	43959	1.92	390	43252	0.90
Komárno	Nitriansky	101712	1.5	343	61268	0.56			
Košice - okolie	Košice	129544	1.2	196	32849	0.60			
Košice I	Košice	67513	1.2	295	39314	0.75			
Košice II	Košice	82288	1.2	41	11109	0.37			
Košice III	Košice	28748	1.2	135	26992	0.50			
Košice IV	Košice	60126	1.2	487	80426	0.61			
Krupina	Banskobystrický	22182	1.4	66	13388	0.49			
Kysucké Nové Mesto	Žilinský	32914	1.6	384	20605	1.86	177	20491	0.86
Levice	Nitriansky	110824	1.4	375	70155	0.53			
Levoča	Prešovský	33702	1.0	373	18344	2.03	172	17747	0.97
Liptovský Mikuláš	Žilinský	72260	1.2	667	47172	1.41	267	46827	0.57
Lučenec	Banskobystrický	73466	1.0	213	40655	0.52			
Malacky	Bratislavský	74323	1.3	285	54657	0.52			
Martin	Žilinský	96338	1.5	771	56533	1.36	381	57513	0.66
Medzilaborce	Prešovský	11842	1.1	91	6980	1.30	17	6142	0.28
Michalovce	Košický	110705	1.0	512	58929	0.87	211	62790	0.34
Myjava	Trenčiansky	26356	0.9	249	17753	1.40	68	18599	0.37
Námestovo	Žilinský	62664	0.9	668	37029	1.80	207	37659	0.55
Nitra	Nitriansky	161560	1.3	674	99175	0.68			
Nové Mesto nad Váhom	Trenčiansky	62554	1.5	363	40829	0.89	198	46269	0.43
Nové Zámky	Nitriansky	139004	1.3	478	79234	0.60			
Partizánske	Trenčiansky	45596	1.5	494	26492	1.86	186	27585	0.67
Pezinok	Bratislavský	65145	1.3	240	45801	0.52			
Piešťany	Trnavský	62802	1.3	183	40122	0.46			
Poltár	Banskobystrický	21471	2.0	71	12455	0.57			
Poprad	Prešovský	104914	1.4	1059	59072	1.79	364	58098	0.63
Považská Bystrica	Trenčiansky	62438	1.4	505	37822	1.34	343	36092	0.95
Prešov	Prešovský	175610	1.0	724	84781	0.85	472	108271	0.44
Prievidza	Trenčiansky	133980	1.3	1497	76457	1.96	576	77170	0.75
Púchov	Trenčiansky	44310	1.3	782	29455	2.65	461	28017	1.65
Revúca	Banskobystrický	39636	1.7	58	21419	0.27			
Rimavská Sobota	Banskobystrický	84159	1.7	197	46872	0.42			
Rožňava	Košický	62208	1.2	100	34307	0.29			
Ružomberok	Žilinský	56702	1.6	682	34000	2.01	236	33056	0.71
Sabinov	Prešovský	60518	1.4	804	35366	2.27	295	34757	0.85
Šaľa	Nitriansky	51685	1.2	199	31993	0.62			
Senec	Bratislavský	89832	1.4	314	66052	0.48			
Senica	Trnavský	60446	1.2	384	40675	0.94	194	46000	0.42
Skalica	Trnavský	47104	1.2	368	29223	1.26	168	31200	0.54
Snina	Prešovský	36240	1.3	345	19122	1.80	111	19396	0.57
Sobrance	Košický	22819	0.9	135	12986	1.04	43	12966	0.33
Spišská Nová Ves	Košický	99765	1.3	739	54279	1.36	361	53712	0.67
Stará Ľubovňa	Prešovský	53954	1.2	805	28749	2.80	354	27234	1.30
Stropkov	Prešovský	20532	1.1	125	10494	1.19	63	10764	0.59
Svidník	Prešovský	32564	1.1	220	16631	1.32	85	16705	0.51
Topoľčany	Nitriansky	70132	1.4	748	44627	1.68	330	50253	0.66
Trebišov	Košický	105353	0.9	400	68503	0.58			
Trenčín	Trenčiansky	114523	1.2	832	73424	1.13	434	72546	0.60
Trnava	Trnavský	132454	1.2	557	92215	0.60			
Turčianske Teplice	Žilinský	15884	1.7	112	11287	0.99	54	12210	0.44
Tvrdošín	Žilinský	36180	1.3	369	18541	1.99	164	20502	0.80
Veľký Krtíš	Banskobystrický	43473	1.2	76	24652	0.31			
Vranov nad Topľou	Prešovský	80766	1.4	460	43552	1.06	281	45424	0.62
Žarnovica	Banskobystrický	26152	1.4	105	16272	0.65			
Žiar nad Hronom	Banskobystrický	46862	0.8	108	26260	0.41			
Žilina	Žilinský	158043	1.5	1392	111155	1.25	512	103898	0.49
Zlaté Moravce	Nitriansky	40572	0.9	156	26180	0.60			
Zvolen	Banskobystrický	68758	1.4	276	39422	0.70	136	47764	0.28

medRxiv preprint doi: https://doi.org/10.1101/2020.12.02.20240648; this version posted December 4, 2020. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.
It is made available under a CC-BY 4.0 International license .

Figure 1: Overview of interventions and pre mass testing epidemiology. Top panel: description of timing and extent of national contact restriction in Slovakia (color intensity indicates intensity of the measures) and timing and extent of the mass testing campaigns. Dots and lines in respective colors show the start and duration of the contact restrictions and the blue dots show the days on which mass testing was conducted, though the highest turnout was usually on the first day. The additional box illustrates contact reducing measures for test positives and those who did chose not to get tested. Bottom panel: SARS-CoV-2 infection incidence as reported by the Slovak Ministry of Health and collected through passive symptom triggered PCR testing. Using the same color coding as in the top panel contact interventions are displayed by horizontal and mass testing campaigns by vertical lines. Data following the respective first mass testing campaign is omitted as mass testing is likely to have interfered with passive surveillance.



medRxiv preprint doi: https://doi.org/10.1101/2020.12.02.20240648; this version posted December 4, 2020. The copyright holder for this preprint **(which was not certified by peer review)** is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.
It is made available under a CC-BY 4.0 International license .

Figure 2: The change in test positivity between mass testing campaigns. Panel A: change in test positivity (1 - cPR) observed from mass testing round 1 to round 2 in the 45 counties that were eligible for both rounds of mass testing. Counties are grouped and color coded into regions. The crude pooled estimate and its 95% confidence bounds are shown as red vertical lines. Panel B: change in test positivity (1 - cPR) observed from the pilot mass testing round to either the first (green) or the second (orange) national round and from the first to the second mass testing round (blue) in the 4 counties that were included in the pilot. Panel C and D: county level test positivity in the first (C) and second (D) round of mass testing. Grey areas indicate counties that were not part of the second round because their test positivity rate was less than 7 per 1000 and hence have no estimates.



medRxiv preprint doi: https://doi.org/10.1101/2020.12.02.20240648; this version posted December 4, 2020. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.
It is made available under a CC-BY 4.0 International license .

Figure 3: Simulated relative effectiveness of the lockdown and the mass testing. Top panel: the change in prevalence of infectious non-quarantining individuals between 10 and 65 years of age as predicted by the microsimulation model. For comparison the observed test-positivity rate is shown in blue. The facets show changes from the pilot to the first round of mass testing (left) and from the pilot to the second round of mass testing (right). Shown scenarios compare the effect of (top to bottom) no additional interventions that limit the growth rate of R_e=1.4, the national lockdown drastically reducing the growth rate to R_e=0.6 and no mass testing being conducted, the national lockdown reducing the growth rate to R_e=1.0 and no mass testing being conducted, no change in growth rate but mass testing, and the national lockdown reducing the growth rate to R_e=1 and mass testing. Bottom panel: Simulated infection incidence of alternative intervention strategies. Simulations are aligned by the date of the first mass test (t=0). The dashed line indicates the timing of the lockdown and the solid lines the timing of the mass testing campaigns. Colors indicate the simulations stratified into whether no mass testing or 1, 2 or 3 testing rounds were performed and the effectiveness of the lockdown measures. Red and yellow dots indicate the prevalence of infectiousness observed among the non-quarantining age-eligible population, corresponding to the scenarios in the top panel.



medRxiv preprint doi: https://doi.org/10.1101/2020.12.02.20240648; this version posted December 4, 2020. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.
It is made available under a CC-BY 4.0 International license .

Supplementary material to "The effectiveness of population-wide screening in reducing SARS-CoV-2 infection prevalence in Slovakia"

by Martin Pavelka, Kevin Van-Zandvoort, Katharine Sherratt, Sam Abbott, Marek Majdan, CMMID COVID-19 working group, Pavol Jarčuška, Marek Krajčí , Stefan Flasche*$, Sebastian Funk*$

Supplementary Tables and Figures

Figure S1: Proportion of positive tests. Test positivity grouped by different mass testing rounds. Given a sufficiently large sample size, one minus test specificity would be the lowest observable proportion of positive test. The absence of apparent clustering of observations at the lower end of the observed range suggests that even lower value could have been observed and test specificity was not a limiting factor.



medRxiv preprint doi: https://doi.org/10.1101/2020.12.02.20240648; this version posted December 4, 2020. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.
It is made available under a CC-BY 4.0 International license .

Figure S2: Simulated relative effectiveness of the lockdown and the mass testing without adherence to quarantine for household members of test-positives. The change in prevalence of infectious non-quarantining individuals between 10 and 65 years of age as predicted by the microsimulation model. For comparison the observed test-positivity rate is shown in light green. The facets show changes from the pilot to the first round of mass testing (top) and from the pilot to the second round of mass testing (bottom). Shown scenarios compare the effect of (top to bottom) no additional interventions that limit the growth rate of R_e=1.4, the national lockdown drastically reducing the growth rate to R_e=0.6 and no mass testing being conducted, no change in growth rate but mass testing, and the national lockdown substantially reducing the growth rate to R_e=1 and mass testing.



Figure S3: Simulated relative effectiveness of the lockdown and the mass testing over time. Simulated infection incidence of alternative intervention strategies. Simulations are aligned by the date of the first mass test (t=0). The dashed line indicates the timing of the lockdown and the solid lines the timing of the mass testing campaigns. Colors indicate the simulations stratified into whether no mass testing or 1, 2 or 3 testing rounds were performed. In the full household compliance facets all household members quarantine for 10 days if a member was tested positive and in the non compliance facet they did not. In Scenario 1 lockdown had no effect on the reproduction number and in Scenario 2 the reproduction number was reduced to 1. The additional grey line in scenario 2 indicates a scenario where no mass testing was done but the reproduction number was reduced to 0.6.



Figure S4: Google mobility index for Slovakia. The change in mobility in comparison to baseline for a number of settings during 2020 in Slovakia. The mobility data is as provided by Google (https://www.google.com/covid19/mobility/).

medRxiv preprint doi: https://doi.org/10.1101/2020.12.02.20240648; this version posted December 4, 2020. The copyright holder for this preprint **(which was not certified by peer review)** is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.
It is made available under a CC-BY 4.0 International license .



Figure S5: Comparing the microsimulation model population to observed structures in Slovakia. Panel A shows the median relative population distribution across all model runs (dark-green) compared to the UNWPP population estimates for Slovakia in 2020 (light-green), by age-group. Panel B shows the median household contact matrix (left; assuming all household members make one contact per day) compared to the synthetic household contact matrix (right), adjusted for UNWPP population size. Panel C shows the median non-household contact matrix (left) compared to the synthetic non-household contact matrix (right), adjusted for lockdown measures and UNWPP population size.



medRxiv preprint doi: https://doi.org/10.1101/2020.12.02.20240648; this version posted December 4, 2020. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.
It is made available under a CC-BY 4.0 International license .

Figure S6: Hospital bed occupancy with COVID-19 patients in Slovakia during the autumn of 2020. Following an increase particularly during October a sharp the abrupt levelling off in the first week of November suggests a sharp decrease in new admissions coinciding with the timing of the mass testing. Data presented are available from the European Centre for Disease Prevention and Control (https://www.ecdc.europa.eu/en/publications-data/download-data-hospital-and-icu-admission-rates-and-current-occupancy-covid-19)



Additional details for the study

Detailed timeline of national SARS-CoV-2infection control measures adopted in Slovakia

Pre - 1 October

- Compulsory face coverings indoors, in enclosed public places and inside mass transport vehicles
- 1000 limit on number of people in aquaparks
- 1000 outdoors and 500 indoors limit on mass gatherings
- Travellers returning from “high risk” countries or regions are requested to take a PCR test after the fifth day of their arrival or remain in quarantine for 10 days
- Shopping hours between 9am and 11am reserved for the elderly

1 October

- Gatherings limited to max 50 people
- Wedding receptions banned

15 October

- Gatherings limited to max 6 people (indoors or outdoors)
- Online schooling for pupils aged 14 years or older
- Compulsory face coverings including outdoors, if within city limits
- Wake receptions banned
- Indoor gastronomy closed
- Theatres and cinemas closed
- Pubs, clubs and bars closed

medRxiv preprint doi: https://doi.org/10.1101/2020.12.02.20240648; this version posted December 4, 2020. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.
It is made available under a CC-BY 4.0 International license .

- Gyms, swimming pools, aquaparks, spas and other wellness and fitness facilities closed
- Church and religious services suspended

24 October - 1 November

- National stay at home order (lockdown) with the following exceptions:
  - travel to and from place of work
  - accompanying children to and from school
  - the first four grades of elementary schools, nurseries and creche stayed open
  - essential travel and activities (i.e. groceries, pharmacy, doctor surgeries, caring for a family dependant, animal husbandry, walking pets within 100 meter distance from home, funerals, post office, bank, insurance company, cleaning services, car repair services, petrol stations)
  - recreational nature walks

2 November

- same restrictions as 15 October with the addition of closing school for pupils aged 10 year or older.

EpiNow2

We used EpiNow2 to backcalculate infection curves in pilot and non-pilot regions. These were converted to infection prevalence using a detection window of 2-6 days after exposure. This allowed us to estimate the infection prevalence of reported cases at the time of mass testing (*p1*) and in the subsequent mass testing round (*p2*). Thus we define the self adjusted prevalence ratio as the crude prevalence ratio observed in the mass testing campaigns adjusted for the predicted change in prevalence if no mass testing or other interventions were conducted:

$$saPR = cPR * p_1/p_2$$

Regression model

We used a quasi Poisson model that was a priori defined by a choice of available covariates that could have plausibly altered the observed impact of the intervention:

$$log(x) = \sum \beta_{0,i} c_i + \beta_1 r + \beta_2 r a_2 + \beta_3 r p_2 + \beta_4 r R_2 + \sum \beta_{5,i} r g_i + log(N) + \epsilon$$

where

x = number of positive tests in each county
N = number of samples
r = round indicator; 0 for first and 1 for second round
c_i = county (categorical)
a_2 = attendance rate of the first national survey (FNS)
p_2 = prevalence observed in the FNS
R_2 = net reproduction number estimated from EpiNow2 for the day of the FNS
g_i = region (categorical)

medRxiv preprint doi: https://doi.org/10.1101/2020.12.02.20240648; this version posted December 4, 2020. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.
It is made available under a CC-BY 4.0 International license .

The model was set up to use the county specific intercept to exactly model the test positivity observed in the first national testing campaign. The round indicator measures the adjusted prevalence ratio (aPR) and the remaining covariates are centered and standardised interaction terms with round to estimate the effects of these variables on the prevalence ratio between the first and second round of mass testing. The number of tests was included as an offset.

Microsimulation model

Model structure

We used an individual-based, probabilistic microsimulation model (IBM) to study the expected reduction in prevalence of (detected) infectiousness under different assumptions.

We up our model to represent an average county of Slovakia .

In our IBM, individuals fall within a age strata (where i is a given age stratum) with relative proportions p_i. They belong to households of mean size m_h (we combine different datasets to simulate a population). The simulation starts when the model population of size N is seeded with at least one SARS-CoV-2 infection, and runs for 365 days.

Births, non-COVID-19 deaths, ageing and migration are omitted from the model given its short timeframe. The study's endpoint of interest is infection, we did not include hospitalisation or clinical outcome status of cases. Infectiousness is assumed to be unaffected by clinical severity, but does differ for asymptomatic, pre-symptomatic and symptomatic cases (see below).

Infection states and transitions

At any time t, individuals within the IBM are within one of the following classes: S (susceptible), E (exposed and latent, i.e. infected but not yet infectious), I_P (infectious but pre-symptomatic), I_C (infectious and symptomatic), I_S (infectious and asymptomatic throughout the infection), or R (removed: recovered and assumed to be immune or deceased). The age-specific probability of becoming a symptomatic case when infected is y_i .

Over any Δt time unit, any given individual has the following binomial probabilities of transitioning to a subsequent state:

$$Pr(S_x \rightarrow E_x) Binomial(1, 1 - e^{-\lambda_{i,x,t}})$$

$$Pr(E_x \rightarrow I_{P,x}) Binomial(1, d_E(t_{E,x}) y_{i,x})$$

$$Pr(E_x \rightarrow I_{S,x}) Binomial(1, d_E(t_{E,x})(1 - y_{i,x}))$$

$$Pr(I_{P,x} \rightarrow I_{C,x}) Binomial(1, d_P(t_{P,x}))$$

$$Pr(I_{C,x} \rightarrow R_x) Binomial(1, d_C(t_{C,x}))$$

$$Pr(I_{S,x} \rightarrow R_x) Binomial(1, d_S(t_{S,x}))$$

where $1 - e^{-\lambda_{i,x,t}}$ is the age-specific instantaneous force of infection experienced by a susceptible individual, as detailed below; and , d_E , d_P , d_C , and d_S are cumulative

medRxiv preprint doi: https://doi.org/10.1101/2020.12.02.20240648; this version posted December 4, 2020. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.
It is made available under a CC-BY 4.0 International license .

distribution functions (CDFs) for the duration of the corresponding states: $d_E(t_{E,x})$ denotes the CDF for the duration of the pre-infectious state evaluated at the time already spent by individual x in that state, and so on.

Transmission dynamics

Over any Δt time unit, susceptible individuals of any age i within each household h move from S to E based on an individual-specific instantaneous force of infection that is the sum of λ due to contacts within the household and λ due to extra-household contacts:

$$\lambda_{i,t,x} = \beta w \frac{I_{P,t,h}+I_{C,t,h}+I_{S,t,h}}{N_{t,h}-1} + \beta \sum_{j=1}^{j=a} U_{ij} \frac{I_{P,t,h'}+I_{C,t,h'}+I_{S,t,h'}}{N_{t,h'}}$$

where β is the probability of infection per contact between a susceptible and infectious person, f is the relative infectiousness of asymptomatic infections, compared to cases that do develop symptoms, w is the mean per-capita intra-household contact rate, assuming random mixing within the household. U is the contact matrix outside the household for the total number of contacts made between individuals aged i with individuals aged j. h denotes individuals within the household itself, while h' denotes individuals in the population excluding the household itself). $I_{P,t}$, $I_{C,t}$, and $I_{S,t}$ represent the total number of infectious individuals not in quarantine at time t.

We assume that all individuals within the household make one contact per day, and calculate the expected population-wide intra-household contact matrix W where W_{ij} is the sum of all aged individuals aged i living together with household members of age j, divided by the model population size aged i. We ensure that the average contact rates are such that the total number of extra-household contacts are symmetric between age-groups, and calculate the population-level contact matrix, $Z = W + U$.

The basic reproduction number R_0 is then defined as the average number of secondary infections generated by a typical infected individual in a fully susceptible population, and is computed as the dominant eigenvalue of the next generation matrix (NGM) of the corresponding compartmental model structure to our IBM model, defined as:

$$NGM_{ij} = \beta Z_{ij}(y_i(\overline{d_P} + \overline{d_C}) + (1 - y_i)f\overline{d_S})$$

where accents indicate the expected (average) values. Lastly, β is the ratio of this eigenvalue and the R_0 value assumed in the simulation (see below).

We validated the calculated R_0 value through this method by running multiple iterations of the model using a different seed for the random number generator, and calculating the average number of secondary cases derived from all infectious individuals who completed their period of infectiousness in the first 30 days of the simulation.

Testing and lockdown

We simulate an epidemic using a timestep of $\Delta t = 1day$. The first round of mass testing is introduced at time t_g when the prevalence of infectiousness in the model reaches a predefined threshold (as observed in the pilot round of mass testing in the county). If introduced, the second and third rounds of testing are introduced on days $t_g + 7$ and $t_g + 14$.

medRxiv preprint doi: https://doi.org/10.1101/2020.12.02.20240648; this version posted December 4, 2020. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.
It is made available under a CC-BY 4.0 International license .

When testing is introduced, we assume that any individual x attends mass-testing with probability z_t. We calculated this probability as $z_t = \frac{N_{attend,t}}{N_{eligible}} \frac{1}{P_{quarantine}}$, where $N_{attend,t}$ is the observed attendance for the test round introduced at time t, $N_{eligible}$ is the total model population size that is eligible for testing (any individual between the ages of 10 and 65), and $P_{quarantine}$ is the proportion of the model population size that is in quarantine at time t.

Individuals already in quarantine do not attend testing. We assume 100% sensitivity to detect an infectious individual (in state I_P, I_S, or I_C), 0% sensitivity to detect an infected but not yet infectious individual (in state E), and 100% specificity for any individual not currently infected. Those who test positive are assumed to comply with quarantine measures with probability C_p, and any of their household members not already quarantining are assumed to comply with probability C_h. We also assume the same probability C_h to quarantine individuals who do not attend mass-testing, but are eligible (between the ages of 10 and 65).

To implement scenarios with lockdown, we first calculated the effective reproduction number in the two weeks before the first round of mass-testing would be implemented, between $t_g - 14$ and t_g. We then started a new model run using the same seed for the random number generator, and implemented a lockdown scenario by changing the value for the probability of effective contact β from the time of implementation of lockdown with $\beta^* = \beta \frac{R_E^*}{R_E}$, where R_E is the estimated effective reproductive number in the period before lockdown and R_E^* is the target value for the effective reproduction number after implementation of lockdown. We assumed the reduced β^* would remain in place for the remainder of the simulation.

Population structure

We simulate a new population within each model iteration by combining estimates for the 2020 Slovak population size, household size by age, and the estimated number of daily contacts made in the household per day.

We simulated a population with target size N by simulating new households until the sum of individuals in all households reached N.

To simulate a household, we randomly sampled i, the age of one individual living in the new household, and drew a value for Y, the household size (ranging from 1 to 6) for those living in the household, from a multinomial distribution where the age-specific distribution of household sizes as estimated in the 2011 Slovak census were used as probabilities for the household size (Eurostat, 2020).

We assumed that normalized age-specific at-home contact rates, W_{ij}^*, calculated as $W_{ij}^* = \frac{W_{ij}}{\sum_{j=1}^{j=a} W_{ij}}$, were proportional to household age distribution (Prem et al, 2020).

We then sampled $Y - 1$ age-groups of household members from a multinomial distribution with age-specific probability of sampling age-group j, $P(j|i) = W_{ij}^* p_j$, where P_j is the probability of sampling any individual from age-group j, following age-specific UNWPP estimates for the population size (UNWPP, 2019).

medRxiv preprint doi: https://doi.org/10.1101/2020.12.02.20240648; this version posted December 4, 2020. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.
It is made available under a CC-BY 4.0 International license .

The median average household size across all modelled populations is 3.7 (3.6-3.7). This is slightly lower than the average household size across all age groups (4.0) as reported in the 2011 Slovak household census (2020, Eurostat - Population by sex, age group, size of household and NUTS 3 regions). Figure S5 compares other key model parameters for the simulated populations with the empirical datasets used. Panel A compares the UNWPP population distribution for Slovakija in 2020 with the median population distribution across all simulated populations. A black area underneath the median population size shows the 95% interval of estimates across all populations, but is not visible in the plot as there is barely any variability across simulated populations, due to the algorithm that was used.

Panel B compares the median household contact matrix across all simulated populations to the synthetic at home contact matrix, where the synthetic matrix has been adjusted with the UNWPP population size estimates to ensure symmetry in the total number of contacts (i.e. total number of contacts of those aged i with j = total number of contacts of those aged j with i). We used the dominant eigenvalue of all matrices to select the matrix representing the median model matrix. The matrices are very similar, though there are slightly less child-adult contacts in the median model matrix compared to the synthetic matrix. The synthetic matrix is generated through extrapolation of contact surveys done in the mid 2000s in other European countries, and may therefore not reflect actual household contact patterns in Slovakia. In addition, the surplus of contacts in the synthetic contact matrix could be due to inclusion of extra-household contacts occurring at the home, which are not included in the model household contact matrix.

Panel C compares the median contact matrix for contacts made outside of the household used in the model, with the contact matrix for non-home contacts in the synthetic matrix for Slovakija (adjusted to represent a change in contact patterns due to Covid-19 interventions). The model contact matrices have been made symmetric for the population distribution used in the model, while the synthetic contact matrix has been made symmetrical for the UNWPP 2020 Slovakija contact matrix, but are otherwise identical. As these population distributions are very similar (Panel A), the contact matrices are as well.

Parameter values

The table below lists all parameter values used in the model

Parameter	Description	Value	Source
H	Number of households	See text	Computed within the model
i, j	Age strata in years (number of age strata =)	0-4, 5-9, 10-14, 15-19, 20-24, 25-29, 30-34, 35-39, 40-44, 45-49, 50-54, 55-59, 60-64, 65-69, 70-74, 75+	n/a
p_i	Proportion of people in each age stratum	Resampled within each model iteration	(UNWPP, 2019)
m_h	Mean household size	Resampled within each model iteration	(Eurostat, 2020)
N	Total population size	78,000	Representative for a typical Slovak county
N_h	Number of people in each household	Resampled within each model iteration	(Eurostat, 2020)
Δt	Time step for discrete-time simulation	1 day	n/a
d_E	Latent period in days	~ gamma(μ = 2.5, k = 4)	
d_P	Duration of pre-symptomatic infectiousness in days	~ gamma(μ = 2.5, k = 4)	
d_C	Duration of symptomatic infectiousness in days	~ gamma(μ = 2.5, k = 4)	

medRxiv preprint doi: https://doi.org/10.1101/2020.12.02.20240648; this version posted December 4, 2020. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.
It is made available under a CC-BY 4.0 International license .

d_S	Duration of asymptomatic infectiousness in days	~ gamma(μ = 5, k = 4)	Assumed to be the same as duration of total infectious period for clinical cases
y_i	Probability of becoming a symptomatic case, if infected, for age group	Age-dependent, as estimated in Davies et al.	(Davies et al, 2020)
R_0	Basic reproduction number	1.5	Assumption, based on EpiNow2 estimates for in time before testing
f	Relative infectiousness of asymptomatic cases	50%	Assumption
w	Within-household per-capita daily contact rate	1	Assumption
W	Age-dependent contact matrix inside the household	Resampled within each model iteration	(Prem, 2020; UNWPP, 2019; Eurostat, 2020)
U	Age-dependent contact matrix outside the household		(Prem, 2020)
β	Probability of transmission per contact with an infectious individual	See text	Computed within the model
z_t	Proportion of people eligible for testing who are tested	As estimated in mass-testing (0.85, 0.78, 0.78)	(Slovakia MOH, 2020)
c_p	Compliance with quarantine for those who test positive	Variable: 0.0, 1.0	Assumption
c_h	Compliance with quarantine for household members of those who test positive	Variable: 0.0, 1.0	Assumption
R_E^*	Target R_E after lockdown	Variable: 0.6, 1.0	Assumption
P_E	Sensitivity of SARS-CoV-2 laboratory test among individuals in latent class	0	Assumption
P_P	Sensitivity of SARS-CoV-2 laboratory test among individuals in pre-symptomatic infectious class	100%	Assumption
P_C	Sensitivity of SARS-CoV-2 laboratory test among individuals in symptomatic infectious class	100%	Assumption
P_S	Sensitivity of SARS-CoV-2 laboratory test among individuals in asymptomatic infectious class	100%	Assumption

Simulations

We ran a total of 15 scenarios and 200 iterations for each:

Scenario	Lockdown effectiveness (R_E^*)	Number of test rounds	Compliance household members (c_h)
1	N/A	0	N/A
2	N/A	1	100%
3	N/A	2	100%
4	N/A	3	100%
5	N/A	1	0%
6	N/A	2	0%
7	N/A	3	0%
8	1	0	N/A
9	1	1	100%
10	1	2	100%
11	1	3	100%
12	1	1	0%
13	1	2	0%
14	1	3	0%
15	0.6	0	N/A